

RESEARCH ARTICLE

Open Access

Comparison of clinical knowledge management capabilities of commercially-available and leading internally-developed electronic health records

Dean F Sittig^{1*}, Adam Wright², Seth Meltzer³, Linas Simonaitis⁴, R Scott Evans⁵, W Paul Nichol⁶, Joan S Ash⁷, Blackford Middleton³

Abstract

Background: We have carried out an extensive qualitative research program focused on the barriers and facilitators to successful adoption and use of various features of advanced, state-of-the-art electronic health records (EHRs) within large, academic, teaching facilities with long-standing EHR research and development programs. We have recently begun investigating smaller, community hospitals and out-patient clinics that rely on commercially-available EHRs. We sought to assess whether the current generation of commercially-available EHRs are capable of providing the clinical knowledge management features, functions, tools, and techniques required to deliver and maintain the clinical decision support (CDS) interventions required to support the recently defined "meaningful use" criteria.

Methods: We developed and fielded a 17-question survey to representatives from nine commercially available EHR vendors and four leading internally developed EHRs. The first part of the survey asked basic questions about the vendor's EHR. The second part asked specifically about the CDS-related system tools and capabilities that each vendor provides. The final section asked about clinical content.

Results: All of the vendors and institutions have multiple modules capable of providing clinical decision support interventions to clinicians. The majority of the systems were capable of performing almost all of the key knowledge management functions we identified.

Conclusion: If these well-designed commercially-available systems are coupled with the other key socio-technical concepts required for safe and effective EHR implementation and use, and organizations have access to implementable clinical knowledge, we expect that the transformation of the healthcare enterprise that so many have predicted, is achievable using commercially-available, state-of-the-art EHRs.

Background

Following passage of the American Recovery and Reinvestment Act of 2009 [1], which provides significant financial incentives over 5 years to eligible hospitals and physicians that implement certified electronic health records (EHR) with clinical decision support (CDS), defined as providing appropriate information to help guide medical decisions at the time and place of care, the pressure on commercial EHR vendors to deliver

full-featured, easy to use systems has increased significantly. Several recent studies have suggested that EHRs without advanced clinical decision support (CDS) features and functions may not provide the promised improvements in patient safety or quality of care [2,3].

Over the past several years, we have carried out an extensive qualitative research program focusing on the barriers and facilitators to successful adoption and use of advanced, state-of-the-art clinical information systems [4,5]. Much of our early work focused on large, academic, teaching facilities with long-standing programs in clinical information system research and development [6]. Over the past several years, we have begun investigating community hospitals

* Correspondence: dean.f.sittig@uth.tmc.edu

¹UTHealth-Memorial Hermann Center for Healthcare Quality & Safety, School of Biomedical Informatics, University of Texas Health Science Center, Houston, TX, USA

Full list of author information is available at the end of the article

and clinics that rely on commercially-available EHR systems [7]. Over these many years, we have seen many different types of EHR systems implemented in many different settings using many different techniques. A result of this work has been development of a new 8-dimension, socio-technical model that we believe helps explain why some implementations go well and others poorly [8]. Clearly, the quality of the technology (i.e., the first dimension of the model), as measured by the features and functions that are available to end-users, plays a major role in any successful implementation. In an attempt to better quantify our qualitative findings that almost all of the EHR systems that we have investigated have nearly the same basic features and functions, as evidenced by the fact that all of them have been CCHIT-approved, we decided to survey a select group of commercial EHR vendors regarding their clinical decision support and knowledge management capabilities.

In order to provide a reference context for the current extent of CDS capabilities, we compared the survey responses of nine commercial vendors with those of four leading internally-developed non-commercial systems. Chaudhry et al. found that approximately 25% of all English-language, peer-reviewed studies that have been used to demonstrate the increases in quality and safety of patient care originated from only four U.S.A healthcare institutions (Intermountain Health Care, Salt Lake City, UT; Veterans Health Administration, Washington, D.C.; Regenstrief Institute, Indianapolis, IN; Brigham & Women's Hospital, Boston, MA). Furthermore, these four institutions provided the majority (54 of 76) of the high-quality literature Chaudhry identified in this area [9]. Chaudhry concluded that studies from these institutions demonstrate that "a multifunctional system can yield real benefits in terms of increased delivery of care based on guidelines (particularly in the domain of preventive health), enhanced monitoring and surveillance activities, reduction of medication errors, and decreased rates of utilization for potentially redundant or inappropriate care." However, they noted that the methods and systems used by these pioneering organizations to obtain these benefits required iterative refinement of locally-developed systems over many years. They, as well as others, have concluded that such long-term, resource intensive, internal health information technology development projects are not a viable option for institutions interested in participating in the recent HITECH stimulus funding which requires significant functionality be in widespread use by 2012 [10].

Therefore, we sought to assess whether the current generation of commercially-available EHRs are capable of providing the technical infrastructure required to deliver and maintain the clinical decision support interventions required to support the recently defined "meaningful use" criteria [11].

Methods

Survey development

We developed a 17-question survey to assess availability of various clinical knowledge management features, functions, tools, and techniques provided by EHRs currently in use. By "clinical knowledge management", we mean the entire process by which clinical knowledge is created, made available, and maintained within an EHR system. This includes the software tools necessary to organize and define knowledge, along with the organizational procedures necessary to manage this knowledge. At the highest level, this requires a systematic and consistent approach to update and maintain the comprehensive set of clinical decision support functions - when a manual ad hoc approach is no longer feasible. A key assumption of this work is that the clinical knowledge management features and functions these systems provide are independent of whether the EHR is designed for the in-patient or out-patient clinical environment. This survey is a quantitative extension of an on-going multi-site, qualitative research project designed to study facilitators and barriers to successful creation, adoption, and dissemination of advanced, clinical decision support to healthcare organizations of all types across the nation [12]. The survey was further informed by the evolving "meaningful use" definition. Finally, the questions were field tested and refined during multiple in-person interviews and demonstrations of existing functionality with EHR vendors at a national EHR vendor conference.

Survey description

The survey was divided into three parts. The first asked basic questions about the electronic health record (EHR) product. For example, we were interested in knowing the name and version of their current product, the different modules (e.g., computer-based provider order entry (CPOE), results review, clinical documentation, etc.) in their system that were capable of providing clinical decision support, how many clinicians were currently using their system nationwide, and what their EHR delivery model was (e.g., on-site server or via a remote hosting solution). This section was designed to assess the capability of the current generation of EHRs to deliver and manage various CDS interventions.

The second part of the survey asked specific questions about the clinical decision support-related system tools and capabilities that each vendor provides. We asked the following questions (the information in italics was designed to clarify each question):

- Does your system allow the collection of structured data within clinician notes [13]? *Structured data can be collected via drop-down lists, from nested menus, or by matching typed free text to a controlled clinical vocabulary. This structured data is then theoretically available*

for use by the CDS logic. We asked those that responded affirmatively, “what controlled vocabularies they use to record the structured clinical findings (e.g., Medications, Laboratory tests, Allergies, Problems/Diagnoses, Procedures)?”

- Does your system support ad-hoc queries (registries) of patient data by end-user [14]? *These queries could be used to support population management, e.g. to identify a group of similar patients taking a medication that was recently recalled, or in need of a preventive health screening exam.* If yes, we asked, “Do you have a replicated database? And “How often is it updated?”

- Do you support the HL-7 InfoButton standard [15]? *The InfoButton provides a mechanism for the user to click on a “hyperlink” from within the EHR and be taken to a relevant place in an online information resource (e.g., Micromedex or CliniGuide).*

- Does your product support the Arden Syntax for Medical Logic Modules (MLMs) [16]? *MLMs are used to share clinical decision support logic that is most often used to create real-time, point-of-care clinical decision support alerts.*

- Does your system provide a clinical knowledge editor so end-users can modify or create their own clinical decision support interventions [17]? *This editor would enable one of your customers to create “if-then” logic. The editor should help them identify specific coded data items within the EHR’s database (e.g., patient_Date_of_Birth, or Weight) and then apply various mathematical (e.g., Greater than) or Boolean (e.g., AND, OR, NOT) operations and if true send a message or alert to the clinician (e.g., elderly patients should not take valium).*

- Do you have a clinical content management system separate from the EHR that allows users to see when their content should be reviewed/updated [18]? *This content management system would allow end-users to browse, sort, or filter the clinical content (a.k.a., the clinical knowledge) that is used within the EHR to create the CDS. In addition, the content management system should contain some “metadata” that further describes the clinical content (e.g., date it was created, author, clinical condition it treats, where it is used in the EHR, etc.). Users should also be able to search or filter content based on this metadata.*

- Do have a reporting capability that allows healthcare organizations to track the effect and/or usage of the CDS content [10]? (Can you measure the response rate for alerts, or order set usage?) *For example, if you had a rule to remind clinicians to order Hemoglobin A1c (HbA1c) tests on all diabetics at least every year, then you would like a reporting capability that could calculate the percentage of all diabetics who actually got an HbA1c test in the past year. Likewise, you might like to know how often this HbA1c reminder has fired in the*

last year and what percentage of the time clinicians “accepted” or followed this reminder.

The final section of the survey asked questions about Clinical Content. Specific questions included:

- How is the CDS content implemented and maintained (e.g., customer is responsible for configuring and maintaining all content to...vendor maintains all content from a central location along with the EHR database)? *We are particularly interested in where the CDS logic/content resides, who maintains and updates the logic, and the processes that are involved in these updates.*

- Who is your medication drug database supplier [20]?

- Do you support any other third party CDS content vendors (e.g. for order sets or alerts) [21]?

- Do you provide clinical content (alert logic, order sets, condition-specific displays) to your clients [22]? *Some vendors call this a “starter-set” or model system that they expect the client to modify before using in their production system. Some vendors provide their clients with charting templates, condition-specific data displays, and even alerts they can use.* If yes, we asked, “what kind of content? How much do you provide, and how and how often do you update this clinical content?”

- Do you support an on-line collaborative knowledge development environment (perhaps using SharePoint or other Web 2.0 technologies) that allows customers to collaborate asynchronously on CDS content [23]?

- Do you have an Internet accessible repository for clients to share their locally generated clinical decision support content with other clients [24]?

Site selection

We selected nine commercially available EHR vendors (three inpatient; six outpatient) from the 2008 list of Certification Commission on Health Information Technology (CCHIT) approved EHR vendors [25]. The vendors were selected because of their size, market position and willingness to cooperate with research. We selected the top four institutions (all of which combined inpatient and outpatient systems) that were responsible for generating approximately 25% of the citations identified by Chaudhry et al. (2006) [9].

Survey distribution

After receiving Institutional Review Board (IRB) approval from the University of Texas ([HSC-SHIS-09-0323](#)), we sent the survey via email to representatives (e.g., product managers or Chief Medical Information Officers) of the nine EHR vendors as well as to colleagues at each of the four institutions with internally-developed systems. In several instances we followed up our email with a teleconference to clarify one or more of our questions, their responses, or both.

Data analysis

All responses were collated, summarized, and are presented anonymously (as agreed upon by all vendors).

Results

Tables 1 and 2 provide information on the EHR vendors and institutions we surveyed along with the location of their headquarters, current version, and number of clinical users of their system.

Table 3 presents the EHR vendor's responses to all of the clinical decision support and knowledge management questions we posed. Vendors requested their identities remain anonymous, therefore we have randomly ordered the columns of table 3. Table 4 presents a similar summary of the responses to all questions from the institutions with the internally-developed systems.

Summary of key findings

All of the vendors and institutions have multiple modules capable of providing clinical decision support interventions to clinicians. All but one of the vendors offer their products remotely (i.e., via Application Service Provider (ASP) models) as well as via on-site servers. In contrast none of the leading internally-developed systems had this capability, although several of them did allow remote access via browsers.

All but two of the vendors and all of the internally-developed systems allow their clients to control the CDS configuration and updating process, although most vendors bundle the clinical content updates with software upgrades. All of the vendors and internally-developed systems stated that they supported the use of a variety of controlled clinical terminologies (e.g., LOINC, SNOMED, ICD-9, Medcin, RxNorm, etc.) to record structured clinical findings, although not all clients make use of these features. All of the vendors and internally-developed systems provided their customers with the ability to perform *ad hoc* queries or patient registry functionality, although only 5/9 vendors and 2/4 internally-developed systems has a real-time, replicated database for these queries. Only 3/9 vendors and 1/4 internally-developed systems

support the HL-7 InfoButton standard, although three other vendors and all the other internally-developed systems have similar, although non-standard, InfoButton-like functionality. Similarly, 3/9 vendors but none of the four internally-developed systems support the Arden Standard for representing medical knowledge.

Seven of the nine vendors and 2/4 of the internally-developed systems provide their customers with a knowledge editor that allows them to create clinical decision support interventions but only 4/9 vendors and 1/4 internally-developed systems store their content in a content management system separate from the application code. Six of the nine vendors and all of the internally-developed systems have the ability to report on the effectiveness of the CDS interventions they deliver, but only 5/9 vendors can report usage statistics of the CDS itself (e.g., alert override rate, or # of times a specific order set was used) whereas all of the internally-developed systems have this capability. All but one of the surveyed vendors and 2/4 of the internally-developed systems reported that they supported First DataBank as their medication-related knowledge base supplier, while 3/9 vendors and 0/4 internally-developed systems reported allowing their customers to utilize multiple commercial medication knowledge base suppliers. Four of the nine vendors and 3/4 of the internally-developed systems reported being able to utilize (i.e., either incorporate into their systems or access via the internet) CDS content from other commercial vendors (e.g., Zynx, Micromedex, UpToDate, etc.). All of the vendors and the internally-developed systems provided their customers with a "starter set" of clinical content (e.g., alerts, order sets, documentation templates, etc.) and at least irregular updates and 6/9 vendors and 2/4 internally-developed systems provide websites (using web 2.0-type functionality) that allow users to share content with others.

Discussion

The vendors in our sample represent a large cross-section of EHRs used in the United States and internationally. We believe that they are a generally representative

Table 1 Overview of vendors surveyed

Vendor	Version	# Users
Allscripts, Chicago, IL	Professional EHR v8	60,000
Cerner Kansas City, MO	Millennium 2007.19	13,000
Eclipsys, Atlanta GA	Sunrise Clinical Manager v5.0sp4	25,000
e-MDs, Austin, TX	Solution Series v6.3	6,000
Epic Systems Verona, WI	Inpatient Summer '09	160,000
GE Healthcare Wauwatosa, WI	Centricity v9.5	"Thousands"
Greenway Medical Technology Carrollton, GA	PrimeSuite 2008	4,000
NextGen, Horsham, PA	EMR v5.5.27	45,000
Spring Medical Systems Houston, TX	SpringCharts EHR v9.7.5	1,500

Table 2 Overview of internally-developed Systems surveyed

Institution	Version	# Users
Intermountain Health Care Salt Lake City, UT	HELP1 and HELP2	20,000
Partners HealthCare Boston, MA	Longitudinal Medical Record (Outpatient) Brigham Integrated Computer System (Inpatient)	32,515
Regenstrief Institute Indianapolis, IN	Gopher CPOE System (Inpatient and Outpatient)	2,500 clinicians entering orders
Veterans Health Administration, Washington, DC	VistA: Computerized Patient Record System V1.0.27.90	172,000

sample of knowledge management capabilities of EHRs, with the possible exception that smaller and niche vendors are not well represented.

Based on the report by Chaudhry, as well as prevailing attitudes in the academic medical informatics community, we expected a large gap in the technical capabilities between the four “best-in-class”, internally-developed

systems as depicted in the scientific literature and commercial EHR vendor’s products. We were pleasantly surprised to discover that all of the vendors we surveyed acknowledged and appreciated the challenges of clinical decision support, and reported that they provided knowledge management capabilities to help their customers manage their clinical knowledge. In fact, with

Table 3 Vendor responses to clinical decision support and knowledge management questions

Questions	EHR: 1	EHR: 2	EHR: 3	EHR: 4	EHR: 5	EHR: 6	EHR: 7	EHR: 8	EHR: 9
CIS modules with CDS?	Y	Y	Y	Y	Y	Y	Y	Y	Y
EHR delivery model?	Onsite or Remote host	Onsite or Remote host	Onsite or Remote host	Onsite or Remote host	On site only	Onsite or Remote host	Onsite or Remote host	Onsite or Remote host	Onsite or Remote host
CDS content implemented, configured, maintained by customer or vendor?	Customer	Vendor	Vendor	Vendor	Vendor	Customer	Vendor	Customer	Vendor
Collect structured data - controlled vocabularies?	Y - multiple	Y - multiple	Y - multiple	Y - multiple	Y - multiple	Y - multiple	Y - multiple	Y - multiple	Y - multiple
Support ad-hoc queries? Live replicating database?	Ye -live db	Yes - db replicated	Yes. No db	Yes.	Yes - Yes	Yes - Yes	Yes - daily update db	Report library - no	Yes-db replicated
HL7 InfoButton?	Yes	Nonstandard	No	No	Nonstandard	Yes	No	Nonstandard	Yes
Arden Syntax for MLM?	Yes	No	No	No	Nonstandard	No	No	Yes	Yes
Clinical knowledge editor for user to create CDS?	Yes	No	Yes	No	Yes	Yes	Yes	Yes	Yes
Content mgmt system separate from EHR?	Yes	No	Yes	No	Yes	No	No	No	Yes
Reporting capability to track effect/usage of CDS?	Yes	Yes.	Yes	No	Yes	Yes	No	N/A	Yes
Drug database supplier?	Multum, FDB	FDB	Proprietary and FDB.	Medi-Span and FDB	FDB, Medi-Span, Lexicomp, Micromedex	FDB	FDB	FDB	Multum MediSource
3rd party CDS content vendors?	Zynx	Multiple	No	No	Multiple	No	No	No	Zynx
Provide clinical content (starter sets). How often and is it updated?	Y- client community	Y- via web	Y-via web	Yes-irregular	Y-1x/year	Y-irregular	Y - irregular	Y-biweekly via web	Y-1x/year
On-line collaborative knowledge development environment (web 2.0)?	Yes, SharePoint	No, listserv	Yes	No	Yes	Yes	Yes	No, listserv	Yes
Internet repository for clients to share CDS?	Yes	No	Yes	No	Yes	Yes	Yes	No	Yes

Table 4 Responses from the internally-developed EHR systems

Questions	Partners	Regenstrief	Intermountain Health Care	Veterans' Health Administration
CIS modules offered (products that have capability to provide decision support)?	Yes, CPOE, results review, laboratory, outpatient clinical documentation, eMAR, Drug-Drug/Allergy/Lab Interaction checking. BCMA	Yes, CPOE, consults from specialists. Documentation: patient visit notes, ADT summaries, problem lists. Links to test results, notes	Yes, Antibiotic, TPN; Results review; RN, RT, Pharmacist charting; Infection & Disease monitoring, ADR triggers, VTE high risk alerts	Yes, Order menus, sets, quick orders. documentation. Templates. Drug-Drug/Allergy/Lab Interaction checking. BCMA
EHR delivery model?	Onsite	Onsite	Onsite (Server and web based)	Onsite (move to regional server)
CDS content implemented/configured/maintained by customer or vendor?	maintained by informaticians. Most content separate from code viewed thru repository	Clinician/Informaticians update terminology and decision support rules as needed	Users responsible for content. Programmers logic. Updated weekly, larger updates quarterly	CDS maintained, implemented locally. National releases with local configuration.
Collect structured data and which controlled vocabularies?	LOINC, RxNorm, SNOMED, ICD-9, CPT, FDB	ICD-9 & CPT codes avail. Most entered via Regenstrief Dict. mapped to standard vocabs.	Mostly coded data, some dictated notes as free text. LOINC, SNOMED, ICD-9	Not from clinician notes, but reminder templates do. ICD-9 for problem list
Support ad-hoc queries? Live replicating database?	Yes. Yes, weekly to monthly depending on data element	Queries done by data analysts on real database, in response to requests from end users.	Yes, but most reports run on EDW from programmers. Have some canned reports monthly	Yes, use replicated DB, varied update frequency; ProClarity data cubes
HL7 InfoButton?	Yes	No	Yes	No, but something similar
Arden Syntax for MLM?	No	No. Gopher Care Rules corresponds to Arden Syntax.	No (some graduate projects)	No
Clinical knowledge editor for user to create CDS?	Yes	Editor used by infomaticians to create CDS	Ability for knowledgeable users to create CDS	Not by end user. Some local configuration by support staff
Content mgmt system separate from EHR?	Yes	No	Online K repository, most review/updates programmers	No. Pilots are ongoing.
Reporting capability to track effect/usage of CDS content?	Yes	User responses prospectively, not retrospective	Yes, from reports in the EDW	Yes, focused on performance measures not CDS usage
Drug database supplier?	FDB	In-house drug terms	FDB	Internal Drug/Allergy files
3rd party CDS content vendors?	Thompson Micromedex, UpToDate	No	MicroMedex through Info-buttons	MicroMedex, UpToDate, etc. at point of care
Provide clinical content (starter sets). What content is included, how and how often and is it updated?	CDS content provided and maintained by the Informatics groups. Updates distributed on an as needed basis.	Alerts, drug-drug. Formulary, price. Order sets. Default drug strengths, frequencies, or instructions. updates mult/yr.	All alert logic is provided, developed, and maintained by institution. weekly and quarterly updates	Local development of reminders, order menus, document templates.
On-line collaborative knowledge development environment web 2.0?	Yes	No	No	Yes, starting SharePoint sites for national groups
Internet repository for clients to share locally generated CDS?	No	No	No	Some networks have web sites to share CDS

respect to the clinical knowledge management capabilities we surveyed, there appears to be little discernable difference between the systems provided by the surveyed EHR vendors and the internally-developed systems. These results mirror our findings from two recent studies that examined the CDS capabilities of commercially-available clinical information systems [26,27]. The vast majority of commercially-available EHR systems

currently on the market in the U.S.A. today report that they provide an adequate technical infrastructure to support the required CDS and knowledge management capabilities.

Despite the strong technical capabilities provided by vendors, however, previous research has shown that simply providing healthcare organizations with CCHIT-compliant, CDS and KM feature-filled electronic health

record applications will not ensure that every organization will be able to realize the process-level improvements in healthcare quality and patient safety that are possible [28].

First, to achieve maximum benefit, organizations will need to select, implement, and maintain the necessary hardware, networking, and operating system software. Without a robust computing infrastructure, the CDS and knowledge management features and functions will not be able to keep up with end-users demands for sub-second response time and 24 hours a day, 7 days a week availability.

Second, organizations will need to develop or purchase, implement, and maintain the clinical content required to provide the clinical decision support interventions required to help guide clinicians. Managing this content is a critical job of either the end-user, or the end-user in conjunction with the vendor, using the knowledge management tools identified in this survey. Ideally, the individuals involved have informatics training (as described below).

Third, EHR vendors will also have to develop easy to learn and use user interfaces that allow users to interact with the CDS interventions in a seamless manner, in the course of the clinical workflow. Failure to integrate the CDS interventions into the clinicians' routine workflow will undoubtedly result in less than optimal usage. Perhaps one of the main reasons that these internally-developed systems have achieved such great success with their CDS interventions is the fact that all of them have been iteratively refined over many years to more closely approximate the clinical workflow of their users. Another reason could be the sense of ownership and responsibility these developers have to make sure the CDS applications are implemented and maintained.

Fourth, to develop, implement, and maintain all of the required clinical content, new kinds of employees capable of interacting with the information technology and clinical personnel will be required. These clinical "informaticians" will be key in the successful CDS implementation and maintenance efforts. Once again, all of the internally-developed systems were designed, developed, and maintained by such a group of committed, highly-trained clinical informaticians. In addition to managing the clinical content, these informaticians also help redesign the clinicians' workflows to make optimal use of the CDS interventions. Further, clinical practice guideline authors, should take note of the clinical content needs of EMRs for CDS, and aim to deliver guidelines in an unambiguous form as near ready for implementation as possible [29].

Fifth, on top of all these other challenges, the organization must review and modify their clinical workflows and communication patterns [30] along with their

internal policies and procedures to reflect newfound capabilities of CDS-enabled EHRs. Failure to make these modifications can result in significantly less than optimal usage of these systems.

Six, the organization must develop the technical and governance infrastructure required to capture, report on, and evaluate the data necessary to measure and monitor both the inner workings of the CDS as well as the effect of the interventions on clinical processes and patient outcomes [31]. Without such careful evaluation, organizations might inadvertently negatively affect the patients' care that they are working to improve [32].

Finally, the local and national bodies responsible for creating and maintaining the rules and regulations that govern the operation of the healthcare facilities and their workers must be updated to take into account the new capabilities of these CDS-enabled EHR systems [33].

Study limitations

The EHR vendor data presented represents only a small sample of the 2008 CCHIT-approved, commercially available EHR systems. Second, the use of self-reported data from the vendors and the internally-developed systems may result in an overstating or an understating of the current capabilities of their systems. When possible we have spoken directly with customers who have installed and are using these systems, as well as, asking for demonstrations of the systems. To date we have not encountered any "inflation" of capabilities. Finally, even though many of these vendors reported a robust suite of CDS and knowledge management features, our experience, along with the literature [34], tells us that not all customers will be able to make use of this advanced CDS functionality and achieve improvements in healthcare quality, safety or reductions in cost [35].

Another limitation is the fact that each of the systems is different, and the survey questions may over-emphasize or under-emphasize the unique capabilities of each system. However, the survey questions needed to be general and somewhat open-ended, to allow us to capture the broad aspects of what is available. After completing this exploratory analysis, future work may involve more specific questions targeted at a limited subset of products, in order to make better comparisons.

Conclusions

Based on a relatively small sample of commercially-available electronic record systems, the clinical knowledge management capabilities of these systems were similar to those of the leading internally-developed systems. The majority of the vendor-supplied systems were capable of performing almost all of the key knowledge management functions we identified. If these well-

designed systems are coupled with the other key socio-technical concepts required for safe and effective EHR implementation and use, we expect that the transformation of healthcare delivery processes that so many have predicted, is just around the corner.

Acknowledgements

We would like to thank Justin Barnes, Emily Bowman, Sarah Corley, Shailja Dixit, Chuck Frederick, Melissa Honour, James Ingram, Jacob Reider, Jack Smyth, and Sean Thomas for helping us complete the vendor surveys. This work was funded in part by NLM Research Grant RO1-LM006942, AHRQ Contract HHSA29020080010 and a contract from the Office of the National Coordinator for Health Information Technology (ONC #10510592). The sponsors were not involved in the design, conduct or analysis of the study, or in the preparation of this manuscript.

Author details

¹UTHealth-Memorial Hermann Center for Healthcare Quality & Safety, School of Biomedical Informatics, University of Texas Health Science Center, Houston, TX, USA. ²Department of Medicine, Brigham & Women's Hospital, Harvard Medical School, Boston, MA, USA. ³Clinical Informatics Research & Development, Partners Healthcare System, Boston, MA, USA. ⁴Regenstrief Institute, Inc.; Indiana University School of Medicine, Indianapolis, IN, USA. ⁵Department of Medical Informatics, Intermountain Health Care, Department of Biomedical Informatics, University of Utah, Salt Lake City, UT, USA. ⁶Patient Care Services, Veterans Health Administration, Seattle WA, USA. ⁷Department of Medical Informatics, Oregon Health & Science University, Portland, OR, USA.

Authors' contributions

DFS and AW, designed, developed, tested and carried out the survey, analyzed the data, and wrote the paper.
SM - Assisted in carrying out the survey, organizing the data, and reviewed many drafts including the final version of the manuscript.
LS - Provided information for the survey and reviewed many drafts including the final version of the manuscript.
RSE - Provided information for the survey and reviewed many drafts including the final version of the manuscript.
WPN - Provided information for the survey and reviewed many drafts including the final version of the manuscript.
JSA - Obtained funding for the study, helped develop the survey questions and reviewed the final version of the manuscript.
BfM - Obtained funding for the study and reviewed many drafts including the final version of the manuscript.
All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 25 November 2010 Accepted: 17 February 2011

Published: 17 February 2011

References

1. American Recovery and Reinvestment Act (ARRA) of 2009. [http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=111_cong_bills&docid=fh1enr.pdf].
2. Linder JA, Ma J, Bates DW, Middleton B, Stafford RS: **Electronic health record use and the quality of ambulatory care in the United States.** *Arch Intern Med* 2007, **167**(13):1400-5.
3. Himmelstein DU, Wright A, Woolhandler S: **Hospital computing and the costs and quality of care: a national study.** *Am J Med* 2010, **123**(1):40-6.
4. Ash JS, Sittig DF, Seshadri V, Dykstra RH, Carpenter JD, Stavri PZ: **Adding insight: a qualitative cross-site study of physician order entry.** *Int J Med Inform* 2005, **74**(7-8):623-8.
5. Ash JS, Sittig DF, Dykstra R, Wright A, McMullen C, Richardson J, Middleton B: **Identifying best practices for clinical decision support and knowledge management in the field.** *Stud Health Technol Inform* 2010, **160**:806-10.
6. Ash JS, Gorman PN, Lavelle M, Stavri PZ, Lyman J, Fournier L, Carpenter J: **Perceptions of physician order entry: results of a cross-site qualitative study.** *Methods Inf Med* 2003, **42**(4):313-23.
7. Guappone KP, Ash JS, Sittig DF: **Field evaluation of commercial Computerized Provider Order Entry systems in community hospitals.** *AMIA Annu Symp Proc* 2008, **6**:263-7.
8. Sittig DF, Singh H: **A New Sociotechnical Model for Studying Health Information Technology in Complex Adaptive Healthcare Systems.** *Qual Saf Health Care* 2010, **19**(Suppl 3):i68-74.
9. Chaudhry B, Wang J, Wu S, et al: **Systematic review: impact of health information technology on quality, efficiency, and costs of medical care.** *Ann Intern Med* 2006, **144**(10):742-52.
10. Blumenthal D, Glaser JP: **Information technology comes to medicine.** *N Engl J Med* 2007, **356**(24):2527-34.
11. Health Information Technology: **Initial Set of Standards, Implementation Specifications, and Certification Criteria for Electronic Health Record Technology; Interim Final Rule.** Federal Register Vol. 75, No. 8/ Wednesday, January 13, 2010/Rules and Regulations.
12. Ash JS, Sittig DF, Guappone KP, Dykstra RH, Richardson J, Wright A, Carpenter J, McMullen C, Shapiro M, Bunce A, Middleton B: **Computerized Clinical Decision Support and Knowledge Management in Community Settings: A Qualitative Study.** *In revision with: JAMIA* 2011.
13. Rosenbloom ST, Miller RA, Johnson KB, Elkin PL, Brown SH: **Interface terminologies: facilitating direct entry of clinical data into electronic health record systems.** *J Am Med Inform Assoc* 2006, **13**(3):277-88.
14. Wright A, McGlinchey EA, Poon EG, Jenter CA, Bates DW, Simon SR: **Ability to generate patient registries among practices with and without electronic health records.** *J Med Internet Res* 2009, **11**(3):e31.
15. HL-7 InfoButton Standard Proposal. [http://www.hl7.org/Library/Committees/dss/infobutton-proposal-2006-01-04.zip].
16. Kim S, Haug PJ, Rocha RA, Choi I: **Modeling the Arden Syntax for medical decisions in XML.** *Int J Med Inform* 2008, **77**(10):650-6.
17. Hripcsak G: **Writing Arden Syntax Medical Logic Modules.** *Comput Biol Med* 1994, **24**(5):331-63.
18. Goldman DS, Colecchi J, Hongsermeier TM, Maviglia SM: **Knowledge management and content integration: a collaborative approach.** *AMIA Annu Symp Proc* 2008, **6**:953.
19. Pearson SA, Moxey A, Robertson J, Hains I, Williamson M, Reeve J, Newby D: **Do computerised clinical decision support systems for prescribing change practice? A systematic review of the literature (1990-2007).** *BMC Health Serv Res* 2009, **9**:154.
20. Kuperman GJ, Reichley RM, Bailey TC: **Using commercial knowledge bases for clinical decision support: opportunities, hurdles, and recommendations.** *J Am Med Inform Assoc* 2006, **13**(4):369-71.
21. **OpenClinical List of Commercial Clinical Decision Support Vendors.**
22. Wright A, Goldberg H, Hongsermeier T, Middleton B: **A description and functional taxonomy of rule-based decision support content at a large integrated delivery network.** *J Am Med Inform Assoc* 2007, **14**(4):489-96.
23. Wright A, Bates DW, Middleton B, Hongsermeier T, Kashyap V, Thomas SM, Sittig DF: **Creating and sharing clinical decision support content with Web 2.0: Issues and examples.** *J Biomed Inform* 2009, **42**(2):334-46.
24. Sittig DF, Wright A, Simonaitis L, Carpenter JD, Allen GO, Doebbeling BN, Sirajuddin AM, Ash JS, Middleton B: **The state of the art in clinical knowledge management: An inventory of tools and techniques.** *Int J Med Informatics* 2010, **79**(1):44-57.
25. **Certification Commission for Health Information Technology.**
26. Wright A, Sittig DF, Ash JS, Sharma S, Pang JE, Middleton B: **Clinical Decision Support Capabilities of Commercially-available Clinical Information Systems.** *J Am Med Inform Assoc* 2009, **16**(5):637-44.
27. Wright A, Sittig DF, Ash JS, Febowitz J, Meltzer S, McMullen C, Guappone K, Carpenter J, Richardson J, Evans RS, Nicol WP, Middleton B: **Development and Evaluation of a Comprehensive Clinical Decision Support Taxonomy: Comparison of Front-End Tools in Commercial and Internally-developed Electronic Health Record Systems.** *J Am Med Inform Assoc* 2011.
28. Sittig DF, Singh H: **Eight Rights of Safe Electronic Health Record Use.** *Journal of American Medical Association* 2009, **302**(10):1111-3.

29. Sittig DF, Wright A, Ash JS, Middleton B: **A set of preliminary standards recommended for achieving a national repository of clinical decision support interventions.** *AMIA Annu Symp Proc* 2009, 614-8, 2009.
30. Campbell EM, Guappone KP, Sittig DF, Dykstra RH, Ash JS: **Computerized provider order entry adoption: implications for clinical workflow.** *J Gen Intern Med* 2009, **24**(1):21-6.
31. Sittig DF, Campbell E, Guappone K, Dykstra R, Ash JS: **Recommendations for monitoring and evaluation of in-patient Computer-based Provider Order Entry systems: results of a Delphi survey.** *AMIA Annu Symp Proc* 2007, 11:671-5.
32. Strom BL, Schinnar R, Abera F, Bilker W, Hennessy S, Leonard CE, Pifer E: **Unintended effects of a computerized physician order entry nearly hard-stop alert to prevent a drug interaction: a randomized controlled trial.** *Arch Intern Med* 2010, **170**(17):1578-83.
33. Blumenthal D, Tavenner M: **The "meaningful use" regulation for electronic health records.** *N Engl J Med* 2010, **363**(6):501-4.
34. Simon SR, Soran CS, Kaushal R, Jenter CA, Volk LA, Burdick E, Cleary PD, Orav EJ, Poon EG, Bates DW: **Physicians' use of key functions in electronic health records from 2005 to 2007: a statewide survey.** *J Am Med Inform Assoc* 2009, **16**(4):465-70.
35. Romano MJ, Stafford RS: **Electronic Health Records and Clinical Decision Support Systems: Impact on National Ambulatory Care Quality.** *Arch Intern Med* 2011.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1472-6947/11/13/prepub>

doi:10.1186/1472-6947-11-13

Cite this article as: Sittig *et al.*: Comparison of clinical knowledge management capabilities of commercially-available and leading internally-developed electronic health records. *BMC Medical Informatics and Decision Making* 2011 11:13.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
biomedcentral.com



[Skip to main page content](#)

National Institutes of Health

U.S. National Library of Medicine

National Center for Biotechnology Information

[NCBI homepage](#)

Close

Account

Logged in as:

username

- [Dashboard](#) (My NCBI)
- [Publications](#) (My Bibliography)
- [Account settings](#)
- [Log out](#)

[Access keys](#) [NCBI Homepage](#) [MyNCBI Homepage](#) [Main Content](#) [Main Navigation](#)

Search:

Search

[Advanced Clipboard](#)

Save Email

...

Save citation to file

Format: Summary (text)

Create file Cancel

Email citation

To:

Subject:

Body:

Format: Summary

Send email Cancel

Add to Collections

- Create a new collection
- Add to an existing collection

Name your collection:

Name must be less than 100 characters

Choose a collection:

Unable to load your collection due to an error

[Please try again](#)

Add Cancel

Add to My Bibliography

- My Bibliography

Unable to load your delegates due to an error

[Please try again](#)

Add Cancel

Full text links

FULL TEXT
JAMA Internal Medicine [Silverchair Information Systems](#)

[Full-text links](#)

Actions

Cite

[Favorites](#)

Share

-
-
-

Page navigation

- [Title & authors](#)
- [Abstract](#)
- [Similar articles](#)
- [Cited by](#)
- [Publication types](#)
- [MeSH terms](#)
- [Grant support](#)
- [LinkOut - more resources](#)

Title & authors

Arch Intern Med

, 167 (13), 1400-5

2007 Jul 9

Electronic Health Record Use and the Quality of Ambulatory Care in the United States

[Jeffrey A Linder](#)¹, [Jun Ma](#), [David W Bates](#), [Blackford Middleton](#), [Randall S Stafford](#)

Affiliations Expand

Affiliation

- ¹ Division of General Medicine, Brigham and Women's Hospital, 1620 Tremont St, BC-3-2X, Boston, MA 02120, USA. jlinder@partners.org
- PMID: **17620534**
- DOI: [10.1001/archinte.167.13.1400](https://doi.org/10.1001/archinte.167.13.1400)

Electronic Health Record Use and the Quality of Ambulatory Care in the United States

Jeffrey A Linder et al. Arch Intern Med. 2007

Show details

Arch Intern Med
, 167 (13), 1400-5

Authors

[Jeffrey A Linder](#)¹, [Jun Ma](#), [David W Bates](#), [Blackford Middleton](#), [Randall S Stafford](#)

Affiliation

- ¹ Division of General Medicine, Brigham and Women's Hospital, 1620 Tremont St, BC-3-2X, Boston, MA 02120, USA. jlinder@partners.org
- PMID: **17620534**
- DOI: [10.1001/archinte.167.13.1400](https://doi.org/10.1001/archinte.167.13.1400)

Full-text links Cite

[Favorites](#)

Abstract

Background: Electronic health records (EHRs) have been proposed as a sustainable solution for improving the quality of medical care. We assessed the association between EHR use, as implemented, and the quality of ambulatory care in a nationally representative survey.

Methods: We performed a retrospective, cross-sectional analysis of visits in the 2003 and 2004 National Ambulatory Medical Care Survey. We examined EHR use throughout the United States and the association of EHR use with 17 ambulatory quality indicators. Performance on quality indicators was defined as the percentage of applicable visits in which patients received recommended care.

Results: Electronic health records were used in 18% (95% confidence interval [CI], 15%-22%) of the estimated 1.8 billion ambulatory visits (95% CI, 1.7-2.0 billion) in the United States in 2003 and 2004. For 14 of the 17 quality indicators, there was no significant difference in performance between visits with vs without EHR use. Categories of these indicators included medical management of common diseases, recommended antibiotic prescribing, preventive counseling, screening tests, and avoiding potentially inappropriate medication prescribing in elderly patients. For 2 quality indicators, visits to medical practices using EHRs had significantly better performance: avoiding benzodiazepine use for patients with depression (91% vs 84%; $P = .01$) and avoiding routine urinalysis during general medical examinations (94% vs 91%; $P = .003$). For 1 quality indicator, visits to practices using EHRs had significantly worse quality: statin prescribing to patients with hypercholesterolemia (33% vs 47%; $P = .01$).

Conclusion: As implemented, EHRs were not associated with better quality ambulatory care.

Similar articles

- [Quality of US Outpatient Care: Temporal Changes and Racial/Ethnic Disparities](#)
J Ma et al. Arch Intern Med 165 (12), 1354-61. 2005. PMID 15983283.
Measurable quality deficits and modest improvements across time call for greater adherence to evidence-based medicine in US ambulatory settings. Although significant raci ...
- [Electronic Health Record Components and the Quality of Care](#)
S Keyhani et al. Med Care 46 (12), 1267-72. Dec 2008. PMID 19300317.
We found no consistent association between blood pressure control, management of chronic conditions, and specific EHR components. Future research focusing on how an EHR i ...
- [Prevalence of the Prescribing of Potentially Inappropriate Medications at Ambulatory Care Visits by Elderly Patients Covered by the Taiwanese National Health Insurance Program](#)
HY Lai et al. Clin Ther 31 (8), 1859-70. Aug 2009. PMID 19808145.
There was a high prevalence of the prescribing of potentially inappropriate medications at ambulatory care visits by elderly patients in Taiwan in 2001-2004.
- [Advancing Evidence-Based Care for Diabetes: Lessons From the Veterans Health Administration](#)
J Kupersmith et al. Health Aff (Millwood) 26 (2), w156-68. Mar-Apr 2007. PMID 17259199. - Review
The Veterans Health Administration (VHA) is a unique laboratory for using the electronic health record (EHR) to transform health care and accelerate discovery. This is pa ...
- [Review: Electronic Health Records and the Reliability and Validity of Quality Measures: A Review of the Literature](#)
KS Chan et al. Med Care Res Rev 67 (5), 503-27. Oct 2010. PMID 20150441. - Review
Previous reviews of research on electronic health record (EHR) data quality have not focused on the needs of quality measurement. The authors reviewed empirical studies o ...

Show more similar articles [See all similar articles](#)

Cited by 130 PubMed Central articles

- [Improving Inpatient Mental Health Medication Safety Through the Process of Obtaining HIMSS Stage 7: A Case Report](#)
H Sulkers et al. JAMIA Open 2 (1), 35-39. 2018. PMID 31984343.
Although electronic health record systems have been implemented in many health settings globally, how organizations can best implement these systems to improve medication ...
- [Trends and Future Direction of the Clinical Decision Support System in Traditional Korean Medicine](#)

HK Sung et al. J Pharmacopuncture 22 (4), 260-268. Dec 2019. PMID 31970024. - Review

In order to implement and utilize the CDSS in a medical service, clinical information must be generated and managed in a standardized form. For this purpose, standardizat ...

- [A Framework for Analysing Learning Health Systems: Are We Removing the Most Impactful Barriers?](#)
S McLachlan et al. Learn Health Syst 3 (4), e10189. 2019. PMID 31641685.
Successful adoption of LHS should take account of the framework proposed in this paper, especially with respect to its focus on removing barriers that have the most impac ...
- [The Association Between Primary Care Physician Compensation and Patterns of Care Delivery, 2012-2015](#)
A Garcia Mosqueira et al. Inquiry 56, 46958019854965. Jan-Dec 2019. PMID 31179800.
As health systems seek to incentivize physicians to deliver high-value care, the relationship between physician compensation and health care delivery is an important know ...
- [Live Usability Testing of Two Complex Clinical Decision Support Tools: Observational Study](#)
S Richardson et al. JMIR Hum Factors 6 (2), e12471. 2019. PMID 30985283.
Live usability testing of these CDS tools generated insights about their role in the patient-provider interaction. CDS may contribute to the interaction by being simultan ...

Show more "Cited by" articles [See all "Cited by" articles](#)

Publication types

- Research Support, U.S. Gov't, P.H.S.

MeSH terms

- Adolescent
- Adult
- Ambulatory Care / standards *
- Cross-Sectional Studies
- Female
- Health Surveys
- Humans
- Male
- Medical Records Systems, Computerized *
- Middle Aged
- Quality of Health Care
- Retrospective Studies
- Smoking Cessation
- United States

Grant support

- [K08 HS014563/HS/AHRQ HHS/United States](#)
- [R01 HS11313/HS/AHRQ HHS/United States](#)

LinkOut - more resources

- **Full Text Sources**

- [Ovid Technologies, Inc.](#)
- [Silverchair Information Systems](#)

- **Medical**

- [ClinicalTrials.gov](#)

Full-text links [x]

FULL TEXT
JAMA Internal Medicine [Silverchair Information Systems](#)

[x]

Citation text

Copy

Download .RIS

Format: AMA

Send To

- [Clipboard](#)
- [Email](#)
- [Save](#)
- [My Bibliography](#)
- [Collections](#)

[x]

Connect

- [Twitter](#)
- [Facebook](#)
- [YouTube](#)
- [LinkedIn](#)
- [GitHub](#)

- [Blog](#)
- [Support Center](#)

National Center for Biotechnology Information

[8600 Rockville Pike Bethesda, MD 20894 USA](#)

- [About us](#)
- [Contact us](#)
- [Policies](#)
- [FOIA](#)

Popular

- [PubMed](#)
- [PubMed Central](#)
- [Bookshelf](#)
- [PubChem](#)
- [Gene](#)
- [BLAST](#)
- [Nucleotide](#)
- [Protein](#)
- [GEO](#)

Resources

- [Literature](#)
- [Health](#)
- [Genomes](#)
- [Genes](#)
- [Proteins](#)
- [Chemicals](#)

Actions

- [Submit](#)
- [Download](#)
- [Learn](#)
- [Develop](#)
- [Analyze](#)
- [Research](#)

[NLM](#) | [NIH](#) | [HHS](#) | [USA.gov](#)

[Feedback](#)

Advanced

Search

Help

Click here to try the **New PubMed!**

An updated version of PubMed is now available. Come see the new improvements to the interface!

Format:
Abstract ▾

Send to ▾

Full text links



AMIA Annu Symp Proc. 2009 Nov 14;2009:614-8.

A set of preliminary standards recommended for achieving a national repository of clinical decision support interventions.

Sittig DF¹, Wright A, Ash JS, Middleton B.

[+ Author information](#)

Abstract

We are investigating the development, implementation and evaluation of clinical decision support (CDS) projects to advance our understanding of how best to incorporate these interventions into the delivery of healthcare. Our overall goal is to explore how the translation of clinical knowledge into CDS and its incorporation into practice can be routinely achieved to improve the quality of healthcare delivered in the U.S.A. Toward this end, we have developed a 7-step model that provides a framework for clinical decision support-related standards that are necessary if we wish to achieve these goals. We believe that if all commercially available EHR systems had the features and functions described in these recommendations, many more healthcare organizations could begin to develop and implement the basic CDS features that are necessary to radically transform the quality, safety, and cost of the current healthcare system.

PMID: 20351928 PMCID: [PMC2815391](#)

[Indexed for MEDLINE] [Free PMC Article](#)

Publication type, MeSH terms, Grant support [+](#)

LinkOut - more resources [+](#)

GETTING STARTED

[NCBI Education](#)
[NCBI Help Manual](#)
[NCBI Handbook](#)
[Training & Tutorials](#)
[Submit Data](#)

RESOURCES

[Chemicals & Bioassays](#)
[Data & Software](#)
[DNA & RNA](#)
[Domains & Structures](#)
[Genes & Expression](#)
[Genetics & Medicine](#)
[Genomes & Maps](#)
[Homology](#)
[Literature](#)
[Proteins](#)
[Sequence Analysis](#)
[Taxonomy](#)
[Variation](#)

POPULAR

[PubMed](#)
[Bookshelf](#)
[PubMed Central](#)
[BLAST](#)
[Nucleotide](#)
[Genome](#)
[SNP](#)
[Gene](#)
[Protein](#)
[PubChem](#)

FEATURED

[Genetic Testing Registry](#)
[GenBank](#)
[Reference Sequences](#)
[Gene Expression Omnibus](#)
[Genome Data Viewer](#)
[Human Genome](#)
[Mouse Genome](#)
[Influenza Virus](#)
[Primer-BLAST](#)
[Sequence Read Archive](#)

NCBI INFORMATION

[About NCBI](#)
[Research at NCBI](#)
[NCBI News & Blog](#)
[NCBI FTP Site](#)
[NCBI on Facebook](#)
[NCBI on Twitter](#)
[NCBI on YouTube](#)
[Privacy Policy](#)

National Center for Biotechnology Information, U.S. National Library of Medicine
8600 Rockville Pike, Bethesda MD, 20894 USA

[Policies and Guidelines](#) | [Contact](#)

